



FucosEXO™

Lyophilized

STORE AT

-20°C



FOR RESEARCH USE ONLY

Instructions for Use

FucosEXO™ Lyophilized 2000 units (G1-FM1-020)
Process 2 mg glycoprotein

DOWNLOAD INSTRUCTIONS FOR USE



www.genovis.com/ifu-G1-FM1

Lyophilized Enzyme for Hydrolysis of α 1-2,3,4 Fucose

FucosEXO is a mix of α -Fucosidases for efficient hydrolysis of α 1-2, α 1-3 and α 1-4-linked fucose residues on native N- and O-glycosylated proteins or free oligosaccharides. FucosEXO hydrolyzes fucose on glycoproteins under native conditions and display a high activity in pH 6.0-8.0.

The enzymes in FucosEXO Lyophilized are derived from *Akkermansia muciniphila* and *Streptococcus oralis*, and expressed in *E. coli*. Both enzymes contain a His-tag and the molecular weights of the enzymes are 87 kDa and 64 kDa.

UNIT DEFINITION

One unit FucosEXO Lyophilized hydrolyzes fucoses from $\geq 90\%$ of 1 μ g glycoprotein (glycoengineered TNFR) when incubated in 20 mM Tris, pH 6.8 at 37°C for 1 hour.

CONTENT AND STORAGE

FucosEXO Lyophilized is supplied lyophilized in TBS, pH 7.6, with no preservatives added.

FucosEXO Lyophilized is shipped cold, and should be stored at -20°C upon arrival. After reconstitution, FucosEXO Lyophilized is stable for at least 1 month at +4-8°C.

FucosEXO Lyophilized is for R&D use only.

QUALITY CONTROL

FucosEXO Lyophilized is tested to meet the specifications and lot-to-lot consistency.

FucosEXO Lyophilized is tested for absence of microbial contamination with blood agar plates, Sabouraud dextrose agar plates and fluid thioglycollate medium.

YOU MIGHT ALSO BE INTERESTED IN

FucosEXO™ Immobilized

Immobilized enzyme mix for hydrolysis of α 1-2,3,4 fucose in spin columns

SialEXO®

Hydrolysis of sialic acids

GalactEXO™

Hydrolysis of β 1-3,4 galactose

GalNAcEXO™

Hydrolysis of α -linked GalNAcs

Preparations

Additional Materials Required

- Reaction buffer: 20mM Tris, pH 6.8.¹

1. FucosEXO Lyophilized displays high activity in buffers with pH 6.0-8.0, and over a wide range of ionic strengths (0-500mM NaCl). Optimization might be required if a buffer other than the recommended reaction buffer is used.

Hydrolysis of α 1-2,3,4 Fucose

Sample Preparation

Prepare the glycoprotein in the reaction buffer. The final glycoprotein concentration in the reaction should be 0.5-5.0mg/ml.

1. Prepare FucosEXO

1.1 Reconstitute FucosEXO in 50 μ l ddH₂O to a concentration of 40 units/ μ l.

2. Add FucosEXO

2.1 Add 1 unit FucosEXO / 1 μ g glycoprotein.²

3. Enzymatic Reaction

3.1 Incubate for 1-2 h at 37°C.

Note: Optimization of enzyme concentrations and incubation time may be needed depending on the substrate.

2. A higher enzyme concentration may increase digestion efficiency of individual glycoproteins. This requires optimization.

USA & Canada

Genovis Inc.

245 First Street, Suite 1800, Cambridge, MA 02142, USA

Phone: 1-855-782-0084 (toll free)

Fax: 1-858-524-3006

EMEA & Asia

Genovis AB

Box 4, SE-24421 Kävlinge, Sweden

Phone: +46 46 10 12 30

Fax: +46 46 12 80 20

support@genovis.com

www.genovis.com



All rights reserved. Genovis products may be covered by one or more patents, trademarks and copyrights owned or controlled by Genovis AB. For more information about commercial rights, please contact the Genovis team at licensing@genovis.com.

Genovis products are intended for research use only. They are not intended to be used for therapeutic or diagnostic purposes in humans or animals.

All goods and services are sold subject to Genovis' General Terms and Conditions of Sale.

© Genovis AB