



GingisKHAN®

Lyophilized

STORE AT

-20°C



FOR RESEARCH USE ONLY

Instructions for Use

GingisKHAN® Lyophilized 2000 units (B0-GKH-020)
Process 2 mg human IgG1

DOWNLOAD INSTRUCTIONS FOR USE



www.genovis.com/ifu-B0-GKH

Lyophilized Enzyme for Above Hinge Digestion of Human IgG1

GingisKHAN (Kgp) is a cysteine protease that digests human IgG1 at a single site above the hinge, generating intact and homogenous Fab and Fc fragments.

GingisKHAN is a lysine-specific protease, and the digestion site on human IgG1 is ...KSCDK / THTCPPC... The single digestion site is a result of the three-dimensional structure of human IgG1, making this lysine exposed for the enzyme. Additional digestion sites at exposed lysines on the Fc may appear if the N-glycans are removed. GingisKHAN requires mildly reducing conditions (2 mM L-cysteine) to be active, and optimal activity is obtained at 37°C and pH 8.0. The reducing agent is provided together with the enzyme.

GingisKHAN is purified from *Porphyromonas gingivalis*.

UNIT DEFINITION

One unit GingisKHAN Lyophilized digests $\geq 95\%$ of 1 μg human IgG1 when incubated in 0.1 M Tris, pH 8.0 at 37°C in presence of 2 mM L-cysteine for 1 hour.

CONTENT AND STORAGE

GingisKHAN Lyophilized contains two components. The product box is shipped cold, and the two components should be stored at -20°C upon arrival.

- **1 vial GingisKHAN Lyophilized** is supplied lyophilized in 100mM Tris, 75mM NaCl, pH 8.0, with no preservatives added. After reconstitution, GingisKHAN Lyophilized is stable for 2 months at +4-8°C.
- **5 vials GingisKHAN Reducing Agent (10x)** is supplied lyophilized, yielding 20mM L-cysteine upon reconstitution (neutral pH). Please note that reconstituted GingisKHAN Reducing Agent cannot be stored – **use it freshly prepared!**

GingisKHAN Lyophilized is for R&D use only.

QUALITY CONTROL

GingisKHAN Lyophilized is tested to meet the specifications and lot-to-lot consistency.

GingisKHAN Lyophilized is tested for absence of microbial contamination with blood agar plates, Sabouraud dextrose agar plates and fluid thioglycollate medium.

YOU MIGHT ALSO BE INTERESTED IN

GingisKHAN® Fab Kit

Lyophilized enzyme and affinity resin for above hinge digestion of human IgG1 and purification of Fab fragments.

FabALACTICA®

Above hinge digestion of human IgG1

FabDELLO™

Above hinge digestion of human IgG1, including hinge-mutated IgG

Preparations

Important Information

- The GingisKHAN Reducing Agent must be prepared and used the same day, it cannot be stored.¹

Additional Materials Required

- Reaction buffer: 0.1 M Tris, pH 8.0.²

1. Upon reconstitution, the GingisKHAN Reducing Agent may appear cloudy. This does not affect the performance. Make sure to mix it thoroughly before using it in the reaction.
2. Other buffers at pH 7.0-8.0 can be used, but optimization is required. NaCl concentrations above 75 mM reduce the enzymatic activity.

Above Hinge Digestion of Human IgG1

Sample Preparation

- Prepare the human IgG1 in the reaction buffer. The final IgG concentration in the reaction should be 0.5-5 mg/ml.
- 1. Prepare GingisKHAN**
 - 1.1 Reconstitute GingisKHAN Lyophilized in 200 μ l ddH₂O to a concentration of 10 units/ μ l.
 - 1.2 Reconstitute GingisKHAN Reducing Agent in 50 μ l ddH₂O and keep on ice.¹ **Note! Use the same day as it is prepared, it cannot be stored.**
 - 2. Add GingisKHAN**
 - 2.1 Add 1 unit GingisKHAN Lyophilized / 1 μ g IgG.
 - 2.2 Add GingisKHAN Reducing Agent to the reaction mixture. Add 1/10 (v/v) to yield 2 mM L-cysteine in the reaction.
 - 3. Enzymatic Reaction**
 - 3.1 Incubate for 1-2 h³ at 37°C.

3. Digestion time may need to be optimized for individual antibodies.

USA & Canada

Genovis Inc.

245 First Street, Suite 1800, Cambridge, MA 02142, USA

Phone: 1-855-782-0084 (toll free)

Fax: 1-858-524-3006

EMEA & Asia

Genovis AB

Box 4, SE-24421 Kävlinge, Sweden

Phone: +46 46 10 12 30

Fax: +46 46 12 80 20

support@genovis.com

www.genovis.com

All rights reserved. Genovis products may be covered by one or more patents, trademarks and copyrights owned or controlled by Genovis AB. For more information about commercial rights, please contact the Genovis team at licensing@genovis.com.

Genovis products are intended for research use only. They are not intended to be used for therapeutic or diagnostic purposes in humans or animals.

All goods and services are sold subject to Genovis' General Terms and Conditions of Sale.

© Genovis AB