



IgMBRAZOR™

IgMBRAZOR™ Lyophilized

STORE AT

-20°C



FOR RESEARCH USE ONLY

Instructions for Use

IgMBRAZOR Lyophilized 1000 units (M0-IM1-010)

Process 1 mg human IgM

DOWNLOAD INSTRUCTIONS FOR USE



www.genovis.com/ifu-M0-IM1

Preparations

ADDITIONAL MATERIALS REQUIRED

- Digestion buffer: PBS (10mM sodium phosphate, 140mM NaCl, 2.7 mM KCl), pH 7.4.¹

SAMPLE PREPARATION

Prepare the IgM in the digestion buffer.

The final protein concentration in the digestion reaction should be 0.5-10 mg/ml.

1. Optimization may be required if buffers other than the recommended are used. The enzyme is active at pH 5.5-9.0 and tolerates up to 300 mM NaCl.

Detailed Protocol

1. Prepare IgMBRAZOR

- 1.1 Reconstitute IgMBRAZOR in 25 μ l ddH₂O to 40 units/ μ l.

2. Add IgMBRAZOR

- 2.1 Add 1 unit IgMBRAZOR/1 μ g IgM.

3. Digestion

- 3.1 Incubate for 30 min at 37°C².

2. The optimal activity is obtained at 37°C. The digestion can be performed at room temperature using an increased incubation time. Optimization is then required.

USA & Canada

Genovis Inc.

245 First Street, Suite 1800, Cambridge, MA 02142, USA

Phone: 1-855-782-0084 (toll free)

Fax: 1-858-524-3006

EMEA & Asia

Genovis AB

Box 790, SE-220 07 Lund, Sweden

Phone: +46 46 10 12 30

Fax: +46 46 12 80 20

support@genovis.com

www.genovis.com

All rights reserved. Genovis products may be covered by one or more patents, trademarks and copyrights owned or controlled by Genovis AB. For more information about commercial rights, please contact the Genovis team at licensing@genovis.com.

Genovis products are intended for research use only. They are not intended to be used for therapeutic or diagnostic purposes in humans or animals.

All goods and services are sold subject to Genovis' General Terms and Conditions of Sale.

© Genovis AB