



SialEXO®

2-3 Lyophilized

STORE AT

-20°C



FOR RESEARCH USE ONLY

Instructions for Use

SialEXO® 2-3 Lyophilized 500 units (G1-SD2-005)
Process 0.5 mg glycoprotein

DOWNLOAD INSTRUCTIONS FOR USE



www.genovis.com/ifu-G1-SD2

Lyophilized Enzyme for Hydrolysis of α 2-3-linked Sialic Acids

SialEXO 2-3 is a sialidase for specific removal of α 2-3-linked sialic acids on O- and N-glycans. SialEXO 2-3 hydrolyzes sialic acids on glycans under native conditions and displays a high activity in a broad pH range, 7.0 to 9.0.

SialEXO 2-3 is derived from *Akkermansia muciniphila* and expressed in *E. coli*. The enzyme contains a His-tag and has a molecular weight of 66 kDa.

UNIT DEFINITION

One unit of SialEXO 2-3 Lyophilized hydrolyzes α 2-3-linked sialic acids from \geq 90% of 1 μ g glycoprotein (fetuin) when incubated in 20 mM Tris pH 7.5 at 37°C for 1 hour.

CONTENT AND STORAGE

SialEXO 2-3 Lyophilized is supplied lyophilized in TBS, pH 7.6, with no preservatives added.

SialEXO 2-3 Lyophilized is shipped at ambient temperature, and should be stored at -20°C upon arrival. After reconstitution, SialEXO 2-3 Lyophilized is stable for at least 1 month at +4-8°C.

SialEXO 2-3 Lyophilized is for R&D use only.

YOU MIGHT ALSO BE INTERESTED IN

SialEXO® Immobilized

Immobilized enzyme mix for hydrolysis of sialic acids in spin columns

OpeRATOR®

O-glycan-specific protein digestion

GlycOCATCH®

Enrichment of O-glycopeptides

OglyZOR®

Hydrolysis of core 1 O-glycans

Preparations

Additional Materials Required

- Reaction buffer: 20mM Tris pH 7.5.¹

1. SialEXO 2-3 displays high activity in buffers at pH 7.0-9.0.

Hydrolysis of α 2-3-linked Sialic Acids

Sample Preparation

Prepare the glycoprotein in the reaction buffer.

The final glycoprotein concentration in the reaction should be 0.1-5 mg/ml.

Prepare the oligosaccharide in the reaction buffer.

The final oligosaccharide concentration in the reaction should be 1-5 pmol/ μ l.

1. Prepare SialEXO 2-3

1.1 Reconstitute SialEXO 2-3 in 25 μ l ddH₂O to a concentration of 20 units/ μ l.

2. Add SialEXO 2-3

2.1 Add 1 unit SialEXO 2-3 / 1 μ g glycoprotein or to 5 pmol oligosaccharides.²

3. Enzymatic Reaction

3.1 Incubate at 37°C for 1 h³.

2. A higher enzyme concentration may increase digestion efficiency of individual glycoproteins or oligosaccharides. This requires optimization.
3. Longer incubation times may be required depending on the substrate.

USA & Canada

Genovis Inc.

245 First Street, Suite 1800, Cambridge, MA 02142, USA

Phone: 1-855-782-0084 (toll free)

Fax: 1-858-524-3006

EMEA & Asia

Genovis AB

Box 4, SE-24421 Kävlinge, Sweden

Phone: +46 46 10 12 30

Fax: +46 46 12 80 20

support@genovis.com

www.genovis.com



All rights reserved. Genovis products may be covered by one or more patents, trademarks and copyrights owned or controlled by Genovis AB. For more information about commercial rights, please contact the Genovis team at licensing@genovis.com.

Genovis products are intended for research use only. They are not intended to be used for therapeutic or diagnostic purposes in humans or animals.

All goods and services are sold subject to Genovis' General Terms and Conditions of Sale.

© Genovis AB